

**ERRATUM****Open Access**

# Erratum to: Early postoperative mortality after simultaneous or staged bilateral primary total hip arthroplasty: an observational register study from the Swedish Hip Arthroplasty Register

Anne Garland<sup>1,2,4\*</sup>, Ola Rolfson<sup>2,3,5</sup>, Göran Garellick<sup>2,3</sup>, Johan Kärrholm<sup>2,3</sup> and Nils P. Hailer<sup>2,4</sup>**Erratum:**

The original version of this article unfortunately contained a mistake in the section entitled “Revision surgery”. The corrected text is given below:

**Corrected text:**

(Page 5: Results - Revision surgery)

Revision surgery of either hip after the second THA surgery was slightly more common in simultaneously operated patients compared to the patients operated with a staged procedure (Simultaneous surgeries 240 [7.1 %], surgeries within 6 months n 495 [5.1 %], surgeries between 7 and 12 months n 828 [5.3 %] and second surgery after >12 months n 3,573 [6.4 %]).

The unadjusted risk of revision for the simultaneously operated patients compared to the patients operated with staged procedure was [HR 1.6, CI 1.3-1.9] (surgeries within 6 months [HR 1.2, CI 1.1-1.4], surgeries between 7 and 12 months [HR 1.1, CI 1.0-1.3] with second surgery after >12 months as the reference group). When adjusting for sex, age, diagnosis and type of prosthesis fixation this difference in the risk estimates disappeared (simultaneous bilateral [HR 1.1, CI 0.9-1.4], surgeries within 6 months [HR 1.1, CI 1.0-1.3], surgeries between 7 and 12 months [HR 1.1, CI 1.0-1.2] with second surgery after >12 months as the reference group).

**Author details**

<sup>1</sup>Department of Orthopaedics, Visby Hospital, Visby, Sweden. <sup>2</sup>Swedish Hip Arthroplasty Register, Gothenburg, Sweden. <sup>3</sup>Department of Orthopaedics, Institute of Clinical Sciences, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. <sup>4</sup>Department of Orthopaedics, Institute of Surgical Sciences, Uppsala University Hospital, Uppsala, Sweden. <sup>5</sup>Harris Orthopaedic Laboratory, Massachusetts General Hospital, Harvard Medical School, Boston, USA.

Received: 11 September 2015 Accepted: 11 September 2015

Published online: 28 September 2015

**Reference**

1. Garland et al. *BMC Musculoskelet Disord.* 2015;16:77. doi:10.1186/s12891-015-0535-0.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [anne.garland@gotland.se](mailto:anne.garland@gotland.se)

<sup>1</sup>Department of Orthopaedics, Visby Hospital, Visby, Sweden

<sup>2</sup>Swedish Hip Arthroplasty Register, Gothenburg, Sweden

Full list of author information is available at the end of the article

